

INTERVENT-Related Scientific Manuscripts and Abstracts (January, 2023)

1. Gordon NF, Scott CB, Levine BD. Comparison of single versus multiple lifestyle interventions: Are the antihypertensive effects of exercise training and diet-induced weight loss additive? *Am J Cardiol* 1997;79:763-767.
2. Gordon NF, Haskell WA. Comprehensive cardiovascular disease risk reduction in a cardiac rehabilitation setting. *Am J Cardiol* 1997;80:69H-73H.
3. Gordon NF. Comprehensive cardiovascular disease risk reduction in the clinical setting. *Coronary Artery Disease* 1998;9:731-735.
4. Gordon NF, Levinrad LI, Salmon RD, Mitchell BS, Alexander LA. Independent and combined effects of the exercise and nutrition components of a community-based primary prevention program. *J Cardiopulm Rehabil Prev* 1998;18:370.
5. Contractor AS, Alexander LA, Blaschko EA, Levinrad LI, Mitchell BS, Salmon RD, Gordon NF. Effect of sequence of lifestyle intervention on select cardiovascular disease risk factors. *Med Sci Sports Exerc* 1999;31:S239.
6. Gordon NF, English CD, Salmon RD, Willoughby MR, Leighton RF, Haskell WL. Clinical effectiveness of three models for comprehensive cardiovascular disease risk reduction in lower risk patients with coronary artery disease. *Circulation* 2000;102:678.
7. Contractor A, English C, Salmon R, Leighton R, Willoughby M, Gordon N. Clinical effectiveness of a community-based cardiovascular risk reduction program (INTERVENT USA) in lower risk patients with coronary artery disease. *J Cardiopulm Rehabil Prev* 2000;20:294.
8. Gordon NF, English CD, Contractor A, Leighton RF, Salmon RD, Lafranchise EF, Widener WG, Franklin BA. Feasibility and clinical effectiveness of a neurologist supervised, nurse case managed stroke risk reduction program. *Circulation* 2000;102:704.
9. Gordon NF, Salmon RD, Mitchell BS, Faircloth GC, Levinrad LI, Salmon S, Saxon WE, Reid KS. Innovative approaches to comprehensive cardiovascular disease risk reduction in clinical and community-based settings. *Current Atherosclerosis Reports* 2001;3:498-506.
10. Badenhop D, Franklin B, Sperling L, Digenio A, Salmon R, Gordon N. Risk factor status on entry into contemporary phase 2 cardiac rehabilitation programs. *Med Sci Sports Exerc* 2001;33:S65.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

11. Gordon N, Franklin B, Sperling L, Badenhop D, Digenio A, Mitchell B, Salmon R. Stage of readiness to change multiple behaviors at entry to a phase 2 cardiac rehabilitation program. *Med Sci Sports Exerc* 2001;33:S126.
12. Bonzheim K, Watson C, Franklin B, Sperling L, Badenhop D, Digenio A, English C, Salmon R, Gordon N. Need for continued cardiovascular disease risk reduction intervention after completion of a contemporary phase 2 cardiac rehabilitation program. *J Cardiopulm Rehabil Prev* 2001;21:321.
13. Mitchell B, Warman S, Pickel S, Ray T, Salmon R, Gordon N. Using discrepancy analysis to assess and improve participant satisfaction in a comprehensive cardiovascular risk reduction program. *J Cardiopulm Rehabil Prev* 2001;21:305.
14. Faircloth C, Warman S, Pickel S, Salmon R, Mitchell B, Franklin B, Gordon N. Benefit of a worksite-based cardiovascular risk reduction program on employee healthcare claims. *J Cardiopulm Rehabil Prev* 2001;21:316.
15. Widener W, Lafranchise F, Franklin B, Leighton R, English C, Salmon R, Gordon N. Clinical effectiveness of a neurologist supervised, nurse case managed stroke risk reduction program in African American versus Caucasian patients. *J Cardiopulm Rehabil Prev* 2001;21:315.
16. Gordon N, English C, Leighton R, Willoughby M, Franklin B, Salmon R. Comparison of a 12-week phase 2 cardiac rehabilitation program and a physician supervised, nurse case managed cardiovascular disease risk reduction program. *J Cardiopulm Rehabil Prev* 2001;21:307.
17. Gordon N, Salmon R, English C, Levinrad I, Leighton R, Franklin B. A low fat/high complex carbohydrate diet is equally beneficial in individuals with and without diabetes when administered as part of a comprehensive cardiovascular disease risk reduction program. *J Cardiopulm Rehabil Prev* 2001;21:325.
18. Franklin B, Bonzheim K, Warren J, Haapaniemi S, Byl N, Ware L, Barnhart S, Gordon N. Effects of phase 2 cardiac rehabilitation participation on patients with abnormal baseline risk factors: Implications for evaluating program effectiveness. *J Cardiopulm Rehabil Prev* 2001;21:307.
19. Pickel S, Warman S, Mitchell B, Ray T, Gordon N. An innovative continuing education program for nurses that addresses personal improvement and professional development. *American Association of Cardiovascular and Pulmonary Rehabilitation Annual Meeting*, 2001.
20. Gordon N, Salmon R, Faircloth C, Levinrad I, Mitchell B, Saxon W, Reid K, Salmon S. Implementation of an innovative community-based heart disease and stroke risk reduction program (INTER_xVENT). *Preventive Med* 2001;33:S15.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

21. Lafranchise F, Widener W, Franklin B, Salmon R, English C, Leighton R, Gordon N. Cardiovascular disease risk factor status of African American versus Caucasian patients referred to a stroke secondary prevention program. *J Stroke Cerebrovasc Dis* 2001;10:199.
22. Lafranchise EF, Widener WG, Franklin BA, Salmon RD, Leighton RF, English CD, Gordon NF. Need for and clinical effectiveness of a neurologist supervised, nurse case managed stroke risk reduction program. *Stroke* 2001;32:367.
23. Franklin BA, Bonzheim K, Warren J, Haapaniemi S, Byl N, Gordon N. Effects of a contemporary, exercise-based rehabilitation and cardiovascular risk reduction program on coronary patients with abnormal baseline risk factors. *CHEST* 2002;122:338-343.
24. Gordon NF, English CD, Contractor AS, Salmon RD, Leighton RF, Franklin BA, Haskell WL. Effectiveness of 3 models for comprehensive cardiovascular disease risk reduction. *Am J Cardiol* 2002;89:1263-1268.
25. Levinrad LI, Salmon RD, English CD, Faircloth GC, Mitchell BS, Saxon WE, Reid KS, Franklin BA, Gordon NF. Clinical effectiveness of a community-based cardiovascular risk reduction program in participants with and without arthritis. *Med Sci Sports Exerc* 2002;34:S159.
26. Arabatzis K, Salmon RD, Pickel S, Adams LC, Kallish IS, Mitchell BS, English CD, Franklin BA, Gordon NF. Clinical effectiveness and reproducibility of a corporate lifestyle management and cardiovascular disease risk reduction program. *Med Sci Sports Exerc* 2002;34:S93.
27. Watson CE, Salmon RD, Arabatzis K, English CD, Mitchell BS, Faircloth GC, Levinrad LI, Saxon WE, Reid KS, Franklin BA, Gordon NF. Clinical effectiveness of a comprehensive cardiovascular risk reduction program: on-site versus telephone/internet delivery. *Med Sci Sports Exerc* 2002;34:S93.
28. Haapaniemi SS, Franklin BA, Sperling LS, Badenhop DT, Salmon RD, English CD, Gordon NF. Clinical effectiveness of a phase 2 cardiac rehabilitation program in participants with and without arthritis. *Med Sci Sports Exerc* 2002;34:S248.
29. Savona T, Salmon R, English CD, Sperling L, Pickel S, Leighton R, Franklin B, Gordon NF. Clinical effectiveness of a community-based cardiovascular risk reduction program in participants with versus without prediabetes. *J Cardiopulm Rehabil Prev* 2002;22:362.
30. Pickel S, Warman S, Mitchell B, Levinrad I, Leighton R, English C, Salmon R, Franklin B, Gordon N. Clinical effectiveness of a community-based cardiovascular risk reduction program in participants with versus without diabetes. *J Cardiopulm Rehabil Prev* 2002;22:354.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

31. Doughty M, Sperling L, Bishop Lindsay K, Salmon R, Mitchell B, Franklin B, Gordon N. Effect of educational status on clinical outcomes in participants in a contemporary phase 2 cardiac rehabilitation program. *J Cardiopulm Rehabil Prev* 2002;22:349.
32. Sperling L, Kallish S, Thiel J, Leighton R, Levinrad I, Salmon R, Franklin B, Gordon N. Clinical effectiveness of a community-based cardiovascular risk reduction program in participants with versus without the metabolic syndrome. *J Cardiopulm Rehabil Prev* 2002; 22:362.
33. Barat C, English C, Pickel S, Warman S, Salmon R, Hall L, Franklin B, Gordon N. Clinical effectiveness of a community-based cardiovascular risk reduction program in African Americans versus Caucasians. *J Cardiopulm Rehabil Prev* 2002;22:355.
34. Haapaniemi S, Franklin B, Badenhop D, Sperling L, Salmon R, Gordon N. Clinical effectiveness of a phase 2 cardiac rehabilitation program in patients with versus without diabetes. *J Cardiopulm Rehabil Prev* 2002;22:362.
35. Franklin B, Haapaniemi S, Salmon R, Mitchell B, Gordon N. Clinical effectiveness of a phase 2 cardiac rehabilitation program in patients with versus without a self-reported history of depression. *J Cardiopulm Rehabil Prev* 2002;22:366.
36. Webb J, Hall L, Salmon R, English C, Franklin B, Gordon N. Effect of perceived health status on clinical outcomes in participants in a contemporary phase 2 cardiac rehabilitation program. *J Cardiopulm Rehabil Prev* 2002;22:355.
37. Gordon N, Salmon R, English C, Saxon W, Leighton R, Dafoe W, Franklin B. A comprehensive cardiovascular disease risk reduction program that includes a low fat/high complex carbohydrate diet is beneficial in individuals with hypertriglyceridemia. *J Cardiopulm Rehabil Prev* 2002;22:355.
38. Maynard T, Sperling L, Franklin B, Hall L, Salmon R, Gordon N. Effect of a phase 2 cardiac rehabilitation program on serum lipids and lipoproteins in patients with versus without known peripheral arterial disease. *J Cardiopulm Rehabil Prev* 2002;22:349.
39. Gordon NF. New methods of delivering secondary preventive services: The promise of the Internet. *J Cardiopulm Rehabil Prev* 2003;23:349-351.
40. Haapaniemi SS, deJong A, Bonzheim KA, Franklin BA, Salmon RD, Gordon NF. Clinical effectiveness of a contemporary cardiac rehabilitation program in patients with the metabolic syndrome. *Med Sci Sports Exerc* 2003;35:S233.
41. Fowler A, Franklin BA, Salmon RD, Gordon NF. Effectiveness of a contemporary cardiac rehabilitation program in patients with a history of heart failure. *Med Sci Sports Exerc* 2003;35:S298.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

42. Franklin BA, Haapaniemi SS, deJong A, Bonzheim KA, Salmon RD, Gordon NF. Clinical effectiveness of a contemporary cardiac rehabilitation program in obese and non-overweight patients. *Med Sci Sports Exerc* 2003;35:S45.
43. Savona T, English CD, Selman MA, Allen K, Blakely J, Rose A, Skiljan R, Salmon R, Franklin B, Gordon N. Clinical effectiveness of a telephone/internet delivered comprehensive cardiovascular risk reduction program in participants from rural communities. *J Cardiopulm Rehabil Prev* 2003;23:371.
44. Salmon R, Selman M, English C, Rose A, Blakely J, Allen K, Skiljan R, Franklin B, Gordon N. Effect of a telephone/internet delivered comprehensive cardiovascular risk reduction program on self-reported health status in participants from rural communities. *J Cardiopulm Rehabil Prev* 2003;23:370.
45. Franklin B, Hall L, Biggerstaff D, Sperling L, Pickel S, Kallish S, Thiel J, Mitchell B, Salmon R, Gordon N. Clinical effectiveness of a comprehensive cardiovascular risk reduction program in female versus male participants. *J Cardiopulm Rehabil Prev* 2003;23:375.
46. Hall L, Franklin B, Biggerstaff D, Pickel S, Sperling L, Thiel J, Kallish S, Mitchell B, Salmon R, Gordon N. Clinical effectiveness of a comprehensive cardiovascular risk reduction program in pre-menopausal versus post-menopausal women. *J Cardiopulm Rehabil Prev* 2003;23:376.
47. Haapaniemi S, Franklin B, Mitchell B, Levinrad I, Saxon W, Reid K, Faircloth C, Salmon R, Gordon N. Effect of educational status on clinical outcomes in participants in a comprehensive lifestyle management and cardiovascular risk reduction program. *J Cardiopulm Rehabil Prev* 2003;23:376.
48. Webb J, Hall L, Franklin B, Sperling L, Lewis A, Faircloth C, Levinrad I, Salmon R, Gordon N. Getting to goal: Is lifestyle intervention worth the effort in patients with an elevated fasting blood glucose level? *J Cardiopulm Rehabil Prev* 2003;23:372.
49. Lewis A, Franklin B, Hall L, Sperling L, Faircloth C, Levinrad I, Salmon R, Gordon N. Getting to goal: Is lifestyle intervention worth the effort in patients with hypertension? *J Cardiopulm Rehabil Prev* 2003;23:373.
50. Black S, Stellmach A, Morrin L. Outcomes of a case-managed home program in a cardiac rehabilitation setting. *J Cardiopulm Rehabil Prev* 2003;23:387.
51. Gordon NF, Salmon RD, Franklin BA, Sperling LS, Hall L, Leighton RF, Haskell WL. Effectiveness of therapeutic lifestyle changes in patients with hypertension, hyperlipidemia, and/or hyperglycemia. *Am J Cardiol* 2004;94:1558-1561.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

52. Gordon NF, Salmon RD, Saxon WE, Reid KS, Faircloth GC, Levinrad I, Mitchell BS, Leighton RF, Sperling LS, Haskell WL, Franklin BA. Getting risk factors to goal: Lifestyle intervention is worth the effort in patients with hypertension, hyperlipidemia, and/or hyperglycemia. *J Am Coll Cardiol* 2004;34:126.
53. McGuckin EE, Franklin BA, Allard-Gould PM, Salmon RD, Mitchell BS, Levinrad LI, Faircloth GC, Gordon NF. Effect of cardiac rehabilitation on self-reported health status in married versus unmarried participants. *Med Sci Sports Exerc* 2004;36:S40.
54. Hendrickson KD, Franklin BA, Allard-Gould PM, Salmon RD, Mitchell BS, Levinrad LI, Faircloth GC, Gordon NF. Clinical effectiveness of a phase 2 cardiac rehabilitation program in married versus unmarried participants. *Med Sci Sports Exerc* 2004;36:S237.
55. DeJong A, Franklin B, Salmon R, Berk M, Gordon N. Effect of a phase 2 cardiac rehabilitation program on self-reported health status in patients with versus without a history of heart failure. *J Cardiopulm Rehabil Prev* 2004;24:343.
56. O'Neal R, Sperling L, Franklin B, Salmon R, Faircloth G, Mitchell B, Levinrad I, Gordon N. Effectiveness of a tobacco cessation intervention, administered as a component of a comprehensive cardiovascular risk reduction program, in participants in different stages of readiness to quit smoking. *J Cardiopulm Rehabil Prev* 2004;24:347.
57. Lewis A, Salmon R, Franklin B, Gordon N. Effect of a phase 2 cardiac rehabilitation program on self-reported health status in obese versus non-overweight patients. *J Cardiopulm Rehabil Prev* 2004;24:348.
58. Draper T, Rubenfire M, Franklin B, Salmon R, Gordon N. Effect of a phase 2 cardiac rehabilitation program on self-reported health status in patients with versus without a self-reported history of depression. *J Cardiopulm Rehabil Prev* 2004;24:355.
59. Gordon NF, Salmon RD, Gordon TL, Biggerstaff ED, Reid KS, Saxon WE, Faircloth GC, Levinrad I, Mitchell BS, Leighton RF, Verma A, Sperling LS, Hall LK, Bairey-Merz CN, Wenger NK, Franklin BA, Haskell WL. Clinical effectiveness of therapeutic lifestyle changes in pre-menopausal versus post-menopausal women. *Circulation* 2005;111:e65.
60. Gordon NF. Innovative approaches to CVD risk reduction: Focus on therapeutic lifestyle changes. *Lipid Spin* 2005;3:7-9.
61. Gordon NF, Salmon RS, Levinrad I, Faircloth GC, Leighton RF, Berk MR, Hall LK, Sperling LS, Dafoe WA, Rubenfire M, Vogel D, Bairey-Merz CN, Haskell WL, Franklin BA. Multicenter study of the clinical effectiveness of a contemporary cardiac rehabilitation program in elderly versus younger patients. *J Am Coll Cardiol* 2005;45:187A.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

62. Gordon NF, Salmon RD, Reid KS, Saxon WE, Faircloth GC, Levinrad I, Mitchell BS, Leighton RF, Verma A, Berk MR, Sperling LS, Haskell WL, Franklin BA. Clinical effectiveness of therapeutic lifestyle changes in African Americans versus Caucasians. *J Am Coll Cardiol* 2005;45:422A.
63. Hendrickson KD, Franklin BA, deJong AT, Salmon RD, Levinrad I, Faircloth GC, Verma A, Leighton RF, Cromwell WC, Gordon NF. Clinical effectiveness of therapeutic lifestyle changes in patients with low-density lipoprotein subclass pattern B versus A. *Med Sci Sports Exerc* 2005;37:S382.
64. deJong AT, Franklin BA, Salmon RD, Levinrad I, Faircloth GC, Wright BS, Reid KS, Saxon WE, Gordon NF. Multicenter study of effect of cardiac rehabilitation on self-reported health status: Elderly versus younger patients. *Med Sci Sports Exerc* 2005;37:S453.
65. Gordon NF, Salmon RD, Levinrad I, Faircloth GC, Wright BS, Reid KS, Saxon WE, Leighton RF, deJong AT, Franklin BA. Clinical effectiveness of a comprehensive lifestyle management and cardiovascular risk reduction program in cancer survivors. *Med Sci Sports Exerc* 2005;37:S312.
66. Fowler AL, Franklin BA, deJong AT, Salmon RD, Levinrad I, Faircloth GC, Wright BS, Reid KS, Saxon WE, Gordon NF. Multi-center study of risk factor status on entry into cardiac rehabilitation: Elderly versus younger patients. *Med Sci Sports Exerc* 2005;37:S227.
67. deJong A, Franklin BA, Salmon RD, Reid KS, Saxon WE, Faircloth GC, Wright BS, Leighton RF, Gordon N. Multi-center study of risk factor status on entry into cardiac rehabilitation: Male versus female patients. *J Cardiopulm Rehabil Prev* 2005;25:297.
68. Sigler M, Salmon RD, Gordon TL, Faircloth GC, Wright BS, Leighton RF, Franklin BA, Gordon N. Effect of gender on clinical responsiveness to therapeutic lifestyle changes. *J Cardiopulm Rehabil Prev* 2005;25:300.
69. Vogel D, Franklin BA, Salmon RD, Reid K, Saxon WE, Faircloth GC, Wright BS, Leighton RF, Gordon N. Multi-center study of risk factor status on completion of a contemporary phase 2 cardiac rehabilitation program: Male versus female patients. *J Cardiopulm Rehabil Prev* 2005;25:304.
70. Gordon N, Draper T, Rubenfire M, Salmon R, Reid K, Saxon W, Faircloth G, Wright B, Leighton R, Franklin B. Prevalence of the metabolic syndrome and its component risk factors on entry into a comprehensive lifestyle management/cardiovascular risk reduction program. *J Cardiopulm Rehabil Prev* 2006;26:260.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

71. Draper T, Rubenfire M, Salmon R, Reid K, Saxon W, Faircloth G, Wright B, Leighton R, Franklin B, Gordon N. Multi-center study of the prevalence of the metabolic syndrome and its component risk factors on entry into a phase 2 cardiac rehabilitation program. *J Cardiopulm Rehabil Prev* 2006; 26:265.
72. Bavikati V, Sperling L, Salmon R, Faircloth G, Leighton R, Franklin B, Gordon N. Clinical effectiveness of therapeutic lifestyle changes in patients with prehypertension. *J Cardiopulm Rehabil Prev* 2006;26:267.
73. Gordon NF and Yamada S. Role of lifestyle intervention in secondary prevention: Challenges, opportunities, and solutions. *Japanese Journal of Cardiac Rehabilitation* 2007;12:10-15.
74. Draper T, Rubenfire M, Salmon R, Reid K, Saxon W, Faircloth G, Wright B, Leighton R, Franklin B, Gordon N. The prevalence of the metabolic syndrome in cardiac rehabilitation patients. *J Am Coll Cardiol* 2007;49:389A.
75. Draper T, Rubenfire M, Smith K, Salmon R, Reid K, Saxon W, Faircloth G, Wright B, Leighton R, Franklin B, Gordon N. Self-reported depression is associated with decreased but considerable benefit from phase 2 cardiac rehabilitation. *J Am Coll Cardiol* 2007;49:390A.
76. Bavikati VV, Sperling LS, Salmon RD, Faircloth GC, Gordon TL, Franklin BA, Gordon NF. Effect of gender and ethnicity on effectiveness of therapeutic lifestyle changes in patients with prehypertension. *J Am Coll Cardiol* 2007;49:392A.
77. Salmon R, Saxon W, Speight T, Drimmer A, Blakely J, Zamora M, Faircloth C, Gordon N. Evaluation of a novel heart rate monitor: comparison with telemetry electrocardiography. *J Cardiopulm Rehabil Prev* 2007;27:267.
78. Salmon RD, Gordon NF, Saxon WE. Effect of gender on effectiveness of therapeutic lifestyle changes in men and women with prediabetes. *Disease Management Association of America Annual Meeting* 2007.
79. Salmon RD, Gordon NF, Reid KS. Gender alters the responsiveness of multiple cardiovascular disease risk factors to a lifestyle management and risk reduction program in participants with prehypertension. *Disease Management Association of America Annual Meeting* 2007.
80. Bavikati VV, Sperling LS, Salmon RD, Faircloth GC, Gordon TL, Franklin BA, Gordon NF. Effect of comprehensive therapeutic lifestyle changes on prehypertension. *Am J Cardiol* 2008;102:1677-1680.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

81. Maron DJ, Forbes BL, Groves JR, Dietrich MS, Sells P, DiGenio. Health-risk appraisal with or without disease management for worksite cardiovascular risk reduction. *Journal of Cardiovascular Nursing* 2008;23:513-518.
82. deJong A, Haapaniemi S, Franklin BA, Case JE, Lewis A, Salmon RD, Gordon NF. Clinical effectiveness of a contemporary cardiac rehabilitation program in myocardial infarction versus bypass surgery patients. *Med Sci Sports Exerc* 2008;40:S329.
83. Haapaniemi S, deJong A, Franklin BA, Case JE, Lewis A, Salmon RD, Gordon NF. Effect of a contemporary cardiac rehabilitation program on self-reported health status: myocardial infarction versus bypass surgery patients. *Med Sci Sports Exerc* 2008;40:S468.
84. Gordon NF, Salmon RD, Alter DA. Effect of a lifestyle health coaching program on multiple cardiovascular disease risk factors in participants with classes I, II and III obesity. *J Cardiopulm Rehabil Prev* 2008;28:280.
85. Gordon NF, Salmon RD, Saxon WE. Effect of lifestyle health coaching on self-reported health status in participants with classes I, II, and III obesity. *Disease Management Association of America Annual Meeting* 2008.
86. Franklin BF, Gordon NF. Contemporary Diagnosis and Management in Cardiovascular Exercise. Handbooks in Healthcare Co. (Newtown, PA), 2009.
87. Gordon NF, Salmon RD, Gordon TL, Alter DA, Franklin BA. Effect of gender on responsiveness of multiple cardiovascular disease risk factors to lifestyle health coaching in adults with prediabetes. *Med Sci Sports Exerc* 2009;41:S286.
88. Franklin BA, Salmon RD, Gordon TL, Alter DA, Gordon NF. Effect of lifestyle health coaching on multiple cardiovascular disease risk factors in overweight and obese participants. *Med Sci Sports Exerc* 2009;41:S285.
89. Gordon NF, Salmon RD, Alter DA, Faircloth GC, Wright BS, Harner J, Wilhide CC, III, Bresnick MB, Franklin BA. Effect of lifestyle health coaching on multiple cardiovascular disease risk factors: Comparison with cardiac rehabilitation. *Med Sci Sports Exerc* 2010;42:S461.
90. Franklin BA, Salmon RD, Alter DA, Faircloth GC, Wright BS, Harner J, Wilhide CC, III, Bresnick MB, Gordon NF. Effect of lifestyle health coaching on the prevalence of metabolic syndrome and its component risk factors. *Med Sci Sports Exerc* 2010;42:S460.
91. Habib A, Salmon R, Bishop KL, Sperling L, Pinzon A, Rubenfire M, Reid K, Faircloth G, Wright B, Drimmer A, Franklin B, Alter D, Gordon N. Effect of cardiac rehabilitation on coronary heart disease risk factors in patients taking antidepressants. *J Cardiopulm Rehabil Prev* 2010;30:274.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

92. Bishop KL, Salmon R, Habib A, Sperling L, Pinzon A, Rubenfire M, Reid K, Faircloth G, Wright B, Drimmer A, Franklin B, Alter D, Gordon N. Effect of cardiac rehabilitation on self-reported health status in patients taking antidepressants. *J Cardiopulm Rehabil Prev* 2010;30:274.
93. Bresnick MG, Alter DA, Kamali K, Salmon RD, Wilhide CC, Harner J, Franklin BA, Gordon NF. Compliance with lifestyle health coaching sessions predicts reduction in modifiable risk factors. *Population Health Management* 2010;13:361.
94. Chimowitz MI, Lynn MJ, Derdeyn CP, et al, for the Stenting and Aggressive Medical Management for Preventing Recurrent Stroke in Intracranial Stenosis Trial Investigators. Stenting versus aggressive medical therapy for intracranial arterial stenosis. *N Engl J Med* 2011;365:993-1003.
95. Turan TN, Lynn MJ, Nizam A, et al, for the SAMMPRIS Trial Investigators. Rationale, design, and implementation of Aggressive Risk Factor Management in the Stenting and Aggressive Medical Management for Prevention of Recurrent Stroke in Intracranial Stenosis (SAMMPRIS) trial. *Circ Cardiovasc Qual Outcomes* 2012;5:e51-e60.
96. Gordon NF, Habib A, Salmon RD, Bishop KL, Drimmer A, Reid KS, Wright BS, Faircloth GC, Gordon TL, Franklin BA, Rubenfire M, Gracik T, Sperling LS. Effect of exercise-based cardiac rehabilitation on multiple atherosclerotic risk factors in patients taking antidepressant medication. *Am J Cardiol* 2013;111:346-351.
97. Gordon NF. Clinical effectiveness of lifestyle management programs: Importance of the class effect paradox. *Current Treatment Options* 2013;15:675-680.
98. Turan TN, Caskey MF, Helwig S, Steiner MR, Wright BS, Montgomery J, Lane B, Lynn MJ, Derdeyn CJ, Fiorella D, Janis S, Chimowitz MI. Participant satisfaction with a lifestyle management program in the SAMMPRIS clinical trial. *Stroke* 2013;44:ATP407.
99. Gordon N, Salmon R, Reid K, Wright B, Faircloth G, Allen K, Thiel J, MA, Savona T, Blakely J, Gordon T. Multi-center study of risk factor status on completion of a contemporary phase 2 cardiac rehabilitation program: Temporal trends. *J Cardiopulm Rehabil Prev* 2013;33:338.
100. Derdeyn CP, Chimowitz MI, Lynn MJ, et al, and the SAMMPRIS Trial Investigators. Aggressive medical treatment with or without stenting in high-risk patients with intracranial artery stenosis (SAMMPRIS): The final results of a randomised trial. *Lancet* 2014;383:333-341.
101. Turan TN, Nizam A, Lynn MJ, Montgomery J, Derdeyn CP, Fiorella D, Lane BF, Janis S, Chimowitz MI. Relationship between compliance with the lifestyle modification program and risk factor control in the stenting and aggressive medical management for preventing recurrent stroke in intracranial stenosis (SAMMPRIS) trial. *Stroke* 2014;45:ATMP105.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

102. Sandesara PB, Lambert CT, Gordon NF, Fletcher GF, Franklin BA, Wenger NK, Sperling L. Cardiac rehabilitation and risk reduction: Time to “rebrand and reinvigorate.” *J Am Coll Cardiol* 2015;65:389-395.
103. Gordon NF, Salmon RD, Wright BS, Faircloth GC, Reid KS, Gordon TL. Clinical effectiveness of lifestyle health coaching: Case study of an evidence-based program. *Am J Lifestyle Med* June 30, 2015, DOI:10.1177/15598276155923512015.
104. Chiu D, Klucznik RP, Turan TN, Lynn MJ, McCane CD, Katz LD, Nizam A, Derdeyn P, Fiorella D, Lane BF, Montgomery J, Janis S, Chimowitz MI. Enrollment volume effect on risk factor control and outcomes in the SAMMPRIS trial. *Neurology* 2015;85:2090-2097.
105. Hannah J, Turan TN; Nizam A, Lynn MJ, Derdeyn CP, Fiorella D, Janis LS, Lane BF, Montgomery J, Chimowitz MI, for the SAMMPRIS Investigators. Type and duration of exercise in the SAMMPRIS trial. *Stroke* 2015;46:AWP381.
106. Chaturvedi S, Howard G, Meschia J. Carotid endarterectomy for asymptomatic stenosis. *JAMA Internal Medicine* 2015;175:1241-1242.
107. Gordon N, Wright B, Salmon R, Faircloth G, Savona T, Gordon T. Participant satisfaction with an evidence-based telephone lifestyle health coaching program. *J Cardiopulm Rehabil Prev* 2015;35:289-290.
108. Schultz W, Gordon N, Salmon R, Eapen D, Sperling L. Statins do not attenuate improvements in fasting plasma glucose and multiple coronary heart disease risk factors during exercise-based cardiac rehabilitation. *J Am Coll Cardiol* 2016;67:1841.
109. Constantinou D, Gradidge P, Salmon R, Schultz W, Sperling L, Franklin B, Gordon N. Gender differences in improvements in atherosclerotic cardiovascular disease risk factors during exercise-based cardiac rehabilitation: Impact of statin therapy. *Br J Sports Med* 2016;50:A70-71.
110. Gordon NF, Salmon RD, Sperling LS, Wright BS, Faircloth GC, Gordon TL, Berk MR, Rubenfire M, Franklin BA. Temporal trends in the achievement of atherosclerotic cardiovascular disease risk factor goals during cardiac rehabilitation. *J Cardiopulmonary Rehabil Prev* 2017;37:11-31.
111. Turan TN, Nizam A, Lynn MJ, Egan BM, Le N-A, Lopes-Virella MF, Hermayer KL, Harrell J, Derdeyn CP, Fiorella D, Janis LS, Lane B, Montgomery J, Chimowitz MI. Relationship between risk factor control and vascular events in the SAMMPRIS trial. *Neurology* 2017;88:379-385.
112. Chaturvedi S, Nahab F. Exercise for stroke prevention: The neglected prescription. *Neurology* 2017;88:4 342-343.

**INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)**

113. Gordon NF, Salmon RD, Wright BS, Faircloth GC, Gordon TL, Hargrett M, Berk MR, Franklin BA, Gulati M. Evaluation of the CardioSmart population health management initiative: Changes in health risks and participant satisfaction. *J Am Coll Cardiol* 2017;69:1836.
114. Howard VJ, Meschia JF, Lal BK, et al. Carotid revascularization and medical management for asymptomatic carotid stenosis: protocol of the CREST-2 clinical trials. *Int J Stroke* 2017;12:770-778.
115. Turan TN, Al Kasab S, Nizam A, Lynn MJ, Harrell J, Derdeyn CP, Fiorella D, Janis LS, Lane BF, Montgomery J, Chimowitz MI; SAMMPRIS Investigators. Relationship between risk factor control and compliance with a lifestyle modification program in the Stenting Aggressive Medical Management for Prevention of Recurrent Stroke in Intracranial Stenosis Trial. *J Stroke Cerebrovasc Dis* 2018;27:801-805.
116. Gordon NF, Salmon RD, Salmon MK, Ponnusamy P. Using digital health technology to promote cardiovascular disease risk reduction in secondary prevention. *Lifestyle Medicine (Third Edition)*, Rippe JM, Ed. CRC Press (Boca Raton, FL), 2019:741-750.
117. Turan TN, Kasab SA, Nizam A, Hannah J, Gordon N, Lynn MJ, Derdeyn CP, Fiorella D, Janis LS, Lane BF, Montgomery J, Chimowitz MI, for the SAMMPRIS Investigators. Type and duration of exercise in the SAMMPRIS trial. *Neurologist* 2019;24:10-12.
118. Harzand A, Vakili AA, Alrohaibani A, Abdelhamid S, Gordon N, Thiel J, Benarroch-Gampel J, Teodorescu V, Minton K, Wenger NK, Rajani R, Shah AJ. Rationale and design of a smartphone-enabled, home-based exercise program in patients with symptomatic peripheral arterial disease: The Smart Step randomized trial. *Clin Cardiol* 2020;43:537-545.
119. Salmon MK, Wright BS, Kridl TL, Faircloth GC, Salmon RD, Gordon NF. Comparative effectiveness of lifestyle intervention on fasting plasma glucose in normal weight vs. overweight/obese adults with prediabetes. *Diabetes* 2020; 69: 273.
120. Turan TN, Voeks JH, Chimowitz MI, Roldan A, LeMatty T, Haley W, Lopes-Virella M, Chaturvedi S, Jones M, Heck D, Howard G, Lal BK, Meschia JF, Brott TG. Rationale, design, and implementation of intensive risk factor treatment in the CREST2 trial. *Stroke* 2020;51:2960-2971.
121. Wongvibulsin S, Habeos EE, Huynh PP, Xun H, Shan R, Rodriguez KP, Wang J, Gandapur YK, Osuji N, Shah LM, Spaulding EM, Hung G, Knowles K, Yang WE, Marvel FA, Levin E, Maron DJ, Gordon NF, Martin SS. Digital interventions for cardiac rehabilitation: A systematic literature review. *JIMR* 2021;23:e18773.

INTERVENT-Related Scientific Manuscripts and Abstracts
(January, 2023)

122. George KS, Salmon MK, Reid KS, Wright BS, Kridl TL, Faircloth GC, Salmon RD, Constantinou D, Gordon NF. Gender does not significantly modulate fasting plasma glucose improvements with lifestyle intervention in prediabetes participants with versus without a family history of diabetes. *Am J Prev Cardiol* 2021;7:100264
123. Salmon MK, Gordon NF, Constantinou D, Reid KS, Wright BS, Kridl TL, Faircloth GC. Comparative effectiveness of lifestyle intervention on fasting plasma glucose in normal weight versus overweight and obese adults with prediabetes. *Am J Lifestyle Med* 2022;16:334-341.
124. Gordon NF, Salmon RD, Salmon MK, Ponnusamy P. Using digital health technology to promote cardiovascular disease risk reduction in secondary prevention. *Lifestyle Medicine* (Fourth Edition), Rippe JM, Ed. CRC Press (Boca Raton, FL), 2023 (In Press).